

---

## Placid subsets

P68087\_en

Setzè Concurs de Programació de la UPC - Semifinal (2018-06-20)

---

You are planning a trip for the  $n$  members of a club. However, some of the members dislike other members. Therefore, you decide to choose a subset  $S$  of members such that:

- Inside  $S$ , noone dislikes anyone.
- There is no  $S'$  such that  $S \subset S'$  and such that  $S'$  fulfils the first property. In other words,  $S$  must be maximal.

Given the information about who dislikes who, can you count the number of such subsets?

### Input

Input consists of several cases, each one with  $n$  followed by  $n$  lines with  $n$  characters each. For  $i \neq j$ , the  $j$ -th character of the  $i$ -th line is 'L' or 'D' depending on whether  $i$  likes or dislikes  $j$ . The diagonal has only dots. Assume  $1 \leq n \leq 20$ .

### Output

For every case, print the number of maximal placid subsets.

#### Sample input

```
2
.D
L.

5
.LDDL
D.LDL
DL.LL
LDD.D
LLLL.

6
.LLLLL
L.LLLL
LL.LLL
DLL.LL
LLDL.L
LLLDD.
```

#### Sample output

```
2
3
4
```

### Problem information

Author : Salvador Roura

Generation : 2024-05-02 21:52:28

© Jutge.org, 2006–2024.

<https://jutge.org>