

---

## Diamonds

P75018\_en

Primer Concurs de Programació de la FME (2004-04-29)

---

A very rich prince has exactly  $n$  diamonds. Each diamond  $1 \leq i \leq n$  has a certain value  $v_i$ . Tradition says that, before getting married, the prince has to give a present of value exactly  $V$  to his princess. The prince wants to give her exactly two of his diamonds, but he does not know how to decide *quickly* if he can do it or not. Can you help to this stupid?

For instance, if  $n = 6$  and the value of the diamonds is  $5, 8, 6, 2, 6, 20$ , then it is possible to give a present of value  $V = 10$  ( $8 + 2$ ) or a present of value  $V = 12$  ( $6 + 6$ ), but it is impossible to give a present of value  $V = 9$ .

### Input

Input consists of several cases. Each case begins with the gift value  $V$  (a natural number between 1 and  $10^8$ ) and the number  $n$  of diamonds (a natural number between 1 and  $10^5$ ) in this order. Then come  $n$  natural numbers between 1 and  $10^8$  indicating the value of each diamond. A case with  $V = n = 0$  marks the end of the input.

### Output

For each case, print a line with “married” or “single” depending on whether the prince can give the present or not.

#### Sample input

```
12 6
5 8 6 2 6 20
9 6
5 8 6 2 6 20
0 0
```

#### Sample output

```
married
single
```

### Problem information

Author : Salvador Roura

Translator : Carlos Molina

Generation : 2024-05-02 23:09:41

© Jutge.org, 2006–2024.

<https://jutge.org>